

# Foundations of Computer Science


## Lecture 14

### Advanced Counting

Sequences with Repetition

Union of Overlapping Sets: Inclusion-Exclusion

Pigeonhole Principle



To count complex objects, construct a sequence of “instructions” that can be used to construct the object uniquely. The number of possible *sequences* of instructions equals the number of possible complex objects.

- ① Sum and product Rules.
- ② Build-up counting:  $\binom{n}{k}$ ,  $n$ -bit sequences with  $k$  1's; goody-bags.
- ③ Counting one set by counting another: bijection.
- ④ Permutations and combinations.
- ⑤ Binomial Theorem.

# Today: Advanced Counting

---

- 1 Sequences with repetition.
  - Anagrams.
- 2 Inclusion-exclusion: extending the sum-rule to overlapping sets.
  - Derangements.
- 3 Pigeonhole principle.
  - Social twins.
  - Subset sums.

# Selecting $k$ from $n$ Distinguishable Objects

---

	no repetition	with repetition
$k$ -sequence		
$k$ -subset		

# Selecting $k$ from $n$ Distinguishable Objects

---

	no repetition	with repetition
$k$ -sequence	$\frac{n!}{(n-k)!}$	
$k$ -subset		

# Selecting $k$ from $n$ Distinguishable Objects

---

	no repetition	with repetition
$k$ -sequence	$\frac{n!}{(n-k)!}$	$n^k$
$k$ -subset		

# Selecting $k$ from $n$ Distinguishable Objects

---

	no repetition	with repetition
$k$ -sequence	$\frac{n!}{(n-k)!}$	$n^k$
$k$ -subset	$\binom{n}{k} = \frac{n!}{k!(n-k)!}$	

# Selecting $k$ from $n$ Distinguishable Objects

---

	no repetition	with repetition
$k$ -sequence	$\frac{n!}{(n-k)!}$	$n^k$
$k$ -subset	$\binom{n}{k} = \frac{n!}{k!(n-k)!}$	$\binom{k+n-1}{n-1}$



# Selecting $k$ from $n$ Distinguishable Objects


	no repetition	with repetition
$k$ -sequence	$\frac{n!}{(n-k)!}$	$n^k$
$k$ -subset	$\binom{n}{k} = \frac{n!}{k!(n-k)!}$	$\binom{k+n-1}{n-1}$
$(k_1, k_2, \dots, k_r)$ -sequence		

$(5, 4, 3)$ -sequence of 5●, 4●, 3●

# Selecting $k$ from $n$ Distinguishable Objects

	no repetition	with repetition
$k$ -sequence	$\frac{n!}{(n-k)!}$	$n^k$
$k$ -subset	$\binom{n}{k} = \frac{n!}{k!(n-k)!}$	$\binom{k+n-1}{n-1}$
$(k_1, k_2, \dots, k_r)$ -sequence		

$(5, 4, 3)$ -sequence of 5●, 4●, 3●




# Selecting $k$ from $n$ Distinguishable Objects

	no repetition	with repetition
$k$ -sequence	$\frac{n!}{(n-k)!}$	$n^k$
$k$ -subset	$\binom{n}{k} = \frac{n!}{k!(n-k)!}$	$\binom{k+n-1}{n-1}$
$(k_1, k_2, \dots, k_r)$ -sequence		

$(5, 4, 3)$ -sequence of 5●, 4●, 3●

Choose slots for ●:  $\binom{12}{5}$  ways



subset of slots used for each type


type - ●

$\{1, 3, 4, 7, 11\}$

# Selecting $k$ from $n$ Distinguishable Objects

	no repetition	with repetition
$k$ -sequence	$\frac{n!}{(n-k)!}$	$n^k$
$k$ -subset	$\binom{n}{k} = \frac{n!}{k!(n-k)!}$	$\binom{k+n-1}{n-1}$
$(k_1, k_2, \dots, k_r)$ -sequence		

$(5, 4, 3)$ -sequence of 5●, 4●, 3●



Choose slots for ●:  $\binom{12}{5}$  ways

Then choose slots for ●:  $\binom{7}{4}$  ways

subset of slots used for each type


type - ●      type - ●

$\{1, 3, 4, 7, 11\}$      $\{2, 5, 6, 10\}$

# Selecting $k$ from $n$ Distinguishable Objects

	no repetition	with repetition
$k$ -sequence	$\frac{n!}{(n-k)!}$	$n^k$
$k$ -subset	$\binom{n}{k} = \frac{n!}{k!(n-k)!}$	$\binom{k+n-1}{n-1}$
$(k_1, k_2, \dots, k_r)$ -sequence		

$(5, 4, 3)$ -sequence of 5●, 4●, 3●



Choose slots for ●:  $\binom{12}{5}$  ways

Then choose slots for ●:  $\binom{7}{4}$  ways

Then choose slots for ●:  $\binom{3}{3}$  ways

subset of slots used for each type


type - ●      type - ●      type - ●

$\{1, 3, 4, 7, 11\}$     $\{2, 5, 6, 10\}$     $\{8, 9, 12\}$

# Selecting $k$ from $n$ Distinguishable Objects

	no repetition	with repetition
$k$ -sequence	$\frac{n!}{(n-k)!}$	$n^k$
$k$ -subset	$\binom{n}{k} = \frac{n!}{k!(n-k)!}$	$\binom{k+n-1}{n-1}$
$(k_1, k_2, \dots, k_r)$ -sequence		

$(5, 4, 3)$ -sequence of 5●, 4●, 3●



subset of slots used for each type

type - ●

type - ●

type - ●

$\{1, 3, 4, 7, 11\}$   $\{2, 5, 6, 10\}$   $\{8, 9, 12\}$

Choose slots for ●:  $\binom{12}{5}$  ways

Then choose slots for ●:  $\binom{7}{4}$  ways

Then choose slots for ●:  $\binom{3}{3}$  ways


Product rule:

$$\binom{12}{5,4,3} = \binom{12}{5} \times \binom{7}{4} \times \binom{3}{3}$$

# Selecting $k$ from $n$ Distinguishable Objects

	no repetition	with repetition
$k$ -sequence	$\frac{n!}{(n-k)!}$	$n^k$
$k$ -subset	$\binom{n}{k} = \frac{n!}{k!(n-k)!}$	$\binom{k+n-1}{n-1}$
$(k_1, k_2, \dots, k_r)$ -sequence		

$(5, 4, 3)$ -sequence of 5●, 4●, 3●



subset of slots used for each type

type - ●

type - ●

type - ●

$\{1, 3, 4, 7, 11\}$   $\{2, 5, 6, 10\}$   $\{8, 9, 12\}$

Choose slots for ●:  $\binom{12}{5}$  ways

Then choose slots for ●:  $\binom{7}{4}$  ways

Then choose slots for ●:  $\binom{3}{3}$  ways


Product rule:

$$\begin{aligned}
 \binom{12}{5,4,3} &= \binom{12}{5} \times \binom{7}{4} \times \binom{3}{3} \\
 &= \frac{12!}{5! \cdot 7!} \times \frac{7!}{4! \cdot 3!} \times \frac{3!}{3! \cdot 0!}
 \end{aligned}$$

# Selecting $k$ from $n$ Distinguishable Objects

	no repetition	with repetition
$k$ -sequence	$\frac{n!}{(n-k)!}$	$n^k$
$k$ -subset	$\binom{n}{k} = \frac{n!}{k!(n-k)!}$	$\binom{k+n-1}{n-1}$
$(k_1, k_2, \dots, k_r)$ -sequence		

$(5, 4, 3)$ -sequence of 5●, 4●, 3●



subset of slots used for each type

type - ●      type - ●      type - ●

$\{1, 3, 4, 7, 11\}$     $\{2, 5, 6, 10\}$     $\{8, 9, 12\}$

Choose slots for ●:  $\binom{12}{5}$  ways

Then choose slots for ●:  $\binom{7}{4}$  ways

Then choose slots for ●:  $\binom{3}{3}$  ways

Product rule:


$$\begin{aligned}
 \binom{12}{5,4,3} &= \binom{12}{5} \times \binom{7}{4} \times \binom{3}{3} \\
 &= \frac{12!}{5! \cdot \cancel{7}!} \times \frac{\cancel{7}!}{4! \cdot \cancel{3}!} \times \frac{\cancel{3}!}{3! \cdot 0!} \\
 &= \frac{12!}{5! \cdot 4! \cdot 3!}
 \end{aligned}$$



# Selecting $k$ from $n$ Distinguishable Objects

	no repetition	with repetition
$k$ -sequence	$\frac{n!}{(n-k)!}$	$n^k$
$k$ -subset	$\binom{n}{k} = \frac{n!}{k!(n-k)!}$	$\binom{k+n-1}{n-1}$
$(k_1, k_2, \dots, k_r)$ -sequence		$\binom{k_1 + \dots + k_r}{k_1, k_2, \dots, k_r} = \frac{k!}{k_1! \cdot k_2! \cdot \dots \cdot k_r!}$

$(5, 4, 3)$ -sequence of 5●, 4●, 3●



subset of slots used for each type

type - ●      type - ●      type - ●

$\{1, 3, 4, 7, 11\}$      $\{2, 5, 6, 10\}$      $\{8, 9, 12\}$

Choose slots for ●:  $\binom{12}{5}$  ways

Then choose slots for ●:  $\binom{7}{4}$  ways

Then choose slots for ●:  $\binom{3}{3}$  ways

Product rule:

$$\begin{aligned}
 \binom{12}{5, 4, 3} &= \binom{12}{5} \times \binom{7}{4} \times \binom{3}{3} \\
 &= \frac{12!}{5! \cdot 7!} \times \frac{7!}{4! \cdot 3!} \times \frac{3!}{3! \cdot 0!} \\
 &= \frac{12!}{5! \cdot 4! \cdot 3!}
 \end{aligned}$$

# Anagrams for AARDVARK

---

A sequence of 8 letters: 3A's, 2R's, 1D, 1V, 1K.

# Anagrams for AARDVARK

---

A sequence of 8 letters: 3A's, 2R's, 1D, 1V, 1K.

Number of such sequences is

$$\binom{8}{3, 2, 1, 1, 1} = \frac{8!}{3! \cdot 2! \cdot 1! \cdot 1! \cdot 1!} = 3360.$$

# Anagrams for AARDVARK

---

A sequence of 8 letters: 3A's, 2R's, 1D, 1V, 1K.

Number of such sequences is

$$\binom{8}{3, 2, 1, 1, 1} = \frac{8!}{3! \cdot 2! \cdot 1! \cdot 1! \cdot 1!} = 3360.$$

**Exercise.** What is the coefficient of  $x^2y^3z^4$  in the expansion of  $(x + y + z)^9$ ?


*[Hint: Sequences of length 9 (why?) with 2 x's, 3 y's and 4 z's.]*

# Extending the Sum Rule to Overlapping Sets

---

$$|A \cup B| = |A| + |B| - |A \cap B|.$$


(Breaks  $A \cup B$  into smaller sets.)



# Extending the Sum Rule to Overlapping Sets

---

$$|A \cup B| = |A| + |B| - |A \cap B|.$$




(Breaks  $A \cup B$  into smaller sets.)

**Example.** How many numbers in  $1, \dots, 10$  are divisible by 2 OR 5.

# Extending the Sum Rule to Overlapping Sets

---

$$|A \cup B| = |A| + |B| - |A \cap B|.$$



(Breaks  $A \cup B$  into smaller sets.)

**Example.** How many numbers in  $1, \dots, 10$  are divisible by 2 OR 5.

$$A = \{\text{numbers divisible by 2}\}.$$


$$|A| = 5.$$

$$(|A| = \lfloor 10/2 \rfloor)$$

# Extending the Sum Rule to Overlapping Sets

---

$$|A \cup B| = |A| + |B| - |A \cap B|.$$



(Breaks  $A \cup B$  into smaller sets.)

**Example.** How many numbers in  $1, \dots, 10$  are divisible by 2 OR 5.

$A = \{\text{numbers divisible by 2}\}.$

$$|A| = 5.$$

$$(|A| = \lfloor 10/2 \rfloor)$$

$B = \{\text{numbers divisible by 5}\}.$

$$|B| = 2.$$


$$(|B| = \lfloor 10/5 \rfloor)$$



# Extending the Sum Rule to Overlapping Sets

---

$$|A \cup B| = |A| + |B| - |A \cap B|.$$



(Breaks  $A \cup B$  into smaller sets.)

**Example.** How many numbers in  $1, \dots, 10$  are divisible by 2 OR 5.

$A = \{\text{numbers divisible by 2}\}.$

$$|A| = 5.$$

$$(|A| = \lfloor 10/2 \rfloor)$$

$B = \{\text{numbers divisible by 5}\}.$

$$|B| = 2.$$

$$(|B| = \lfloor 10/5 \rfloor)$$

$A \cap B = \{\text{numbers divisible by 2 AND 5}\}.$


$$|A \cap B| = 1.$$

$$(|A \cap B| = \lfloor 10/\text{lcm}(2, 5) \rfloor)$$

# Extending the Sum Rule to Overlapping Sets

---

$$|A \cup B| = |A| + |B| - |A \cap B|.$$



(Breaks  $A \cup B$  into smaller sets.)

**Example.** How many numbers in  $1, \dots, 10$  are divisible by 2 OR 5.

$$A = \{\text{numbers divisible by 2}\}. \quad |A| = 5. \quad (|A| = \lfloor 10/2 \rfloor)$$

$$B = \{\text{numbers divisible by 5}\}. \quad |B| = 2. \quad (|B| = \lfloor 10/5 \rfloor)$$

$$A \cap B = \{\text{numbers divisible by 2 AND 5}\}. \quad |A \cap B| = 1. \quad (|A \cap B| = \lfloor 10/\text{lcm}(2, 5) \rfloor)$$

$$\mathbf{A \cup B = \{numbers divisible by 2 or 5\}.}$$


$$|A \cup B| = |A| + |B| - |A \cap B| = 5 + 2 - 1 = 6.$$

# Inclusion-Exclusion

---

$$|A_1 \cup A_2 \cup A_3| = |A_1| + |A_2| + |A_3| - |A_1 \cap A_2| - |A_1 \cap A_3| - |A_2 \cap A_3| + |A_1 \cap A_2 \cap A_3|.$$

*Proof.*




# Inclusion-Exclusion

---

$$|A_1 \cup A_2 \cup A_3| = |A_1| + |A_2| + |A_3| - |A_1 \cap A_2| - |A_1 \cap A_3| - |A_2 \cap A_3| + |A_1 \cap A_2 \cap A_3|.$$

*Proof.* Consider  $x \in A_2 \cap A_3$ . How many times is  $x$  counted?




# Inclusion-Exclusion

---

$$|A_1 \cup A_2 \cup A_3| = |A_1| + |A_2| + |A_3| - |A_1 \cap A_2| - |A_1 \cap A_3| - |A_2 \cap A_3| + |A_1 \cap A_2 \cap A_3|.$$

*Proof.* Consider  $x \in A_2 \cap A_3$ . How many times is  $x$  counted?

$$\begin{array}{ccccccc} |A_1| & + & |A_2| & + & |A_3| & - & |A_1 \cap A_2| & - & |A_1 \cap A_3| & - & |A_2 \cap A_3| & + & |A_1 \cap A_2 \cap A_3| \\ 0 & & +1 & & +1 & & 0 & & 0 & & -1 & & 0 \end{array}$$



# Inclusion-Exclusion


---

$$|A_1 \cup A_2 \cup A_3| = |A_1| + |A_2| + |A_3| - |A_1 \cap A_2| - |A_1 \cap A_3| - |A_2 \cap A_3| + |A_1 \cap A_2 \cap A_3|.$$

*Proof.* Consider  $x \in A_2 \cap A_3$ . How many times is  $x$  counted?

$$\begin{array}{ccccccc} |A_1| & + & |A_2| & + & |A_3| & - & |A_1 \cap A_2| & - & |A_1 \cap A_3| & - & |A_2 \cap A_3| & + & |A_1 \cap A_2 \cap A_3| \\ 0 & & +1 & & +1 & & 0 & & 0 & & -1 & & 0 \end{array}$$

Contribution of  $x$  to sum is  $+1$ . Repeat for each region. ■




# Inclusion-Exclusion

$$|A_1 \cup A_2 \cup A_3| = |A_1| + |A_2| + |A_3| - |A_1 \cap A_2| - |A_1 \cap A_3| - |A_2 \cap A_3| + |A_1 \cap A_2 \cap A_3|.$$

*Proof.* Consider  $x \in A_2 \cap A_3$ . How many times is  $x$  counted?

$$\begin{array}{ccccccc} |A_1| & + & |A_2| & + & |A_3| & - & |A_1 \cap A_2| & - & |A_1 \cap A_3| & - & |A_2 \cap A_3| & + & |A_1 \cap A_2 \cap A_3| \\ 0 & & +1 & & +1 & & 0 & & 0 & & -1 & & 0 \end{array}$$

Contribution of  $x$  to sum is  $+1$ . Repeat for each region. ■



**Example (Derangements).** Give 3 coats to 3 girls so that noone gets their coat. How many ways?


# Inclusion-Exclusion

$$|A_1 \cup A_2 \cup A_3| = |A_1| + |A_2| + |A_3| - |A_1 \cap A_2| - |A_1 \cap A_3| - |A_2 \cap A_3| + |A_1 \cap A_2 \cap A_3|.$$

*Proof.* Consider  $x \in A_2 \cap A_3$ . How many times is  $x$  counted?

$$\begin{array}{ccccccc} |A_1| & + & |A_2| & + & |A_3| & - & |A_1 \cap A_2| & - & |A_1 \cap A_3| & - & |A_2 \cap A_3| & + & |A_1 \cap A_2 \cap A_3| \\ 0 & & +1 & & +1 & & 0 & & 0 & & -1 & & 0 \end{array}$$

Contribution of  $x$  to sum is  $+1$ . Repeat for each region. ■



**Example (Derangements).** Give 3 coats to 3 girls so that noone gets their coat. How many ways?

$A_i = \{\text{girl } i \text{ gets her coat}\}$ .  $|A_i| = 2!$ .

$A_{ij} = \{\text{girls } i \text{ and } j \text{ get their coats}\}$ .  $|A_{ij}| = 1!$ .

$A_{123} = \{\text{girls 1, 2 and 3 get their coats}\}$ .  $|A_{123}| = 1$ .




# Inclusion-Exclusion

$$|A_1 \cup A_2 \cup A_3| = |A_1| + |A_2| + |A_3| - |A_1 \cap A_2| - |A_1 \cap A_3| - |A_2 \cap A_3| + |A_1 \cap A_2 \cap A_3|.$$

*Proof.* Consider  $x \in A_2 \cap A_3$ . How many times is  $x$  counted?

$$\begin{array}{ccccccc} |A_1| & + & |A_2| & + & |A_3| & - & |A_1 \cap A_2| & - & |A_1 \cap A_3| & - & |A_2 \cap A_3| & + & |A_1 \cap A_2 \cap A_3| \\ 0 & & +1 & & +1 & & 0 & & 0 & & -1 & & 0 \end{array}$$

Contribution of  $x$  to sum is  $+1$ . Repeat for each region. ■



**Example (Derangements).** Give 3 coats to 3 girls so that noone gets their coat. How many ways?

$A_i = \{\text{girl } i \text{ gets her coat}\}$ .  $|A_i| = 2!$ .

$A_{ij} = \{\text{girls } i \text{ and } j \text{ get their coats}\}$ .  $|A_{ij}| = 1!$ .

$A_{123} = \{\text{girls 1, 2 and 3 get their coats}\}$ .  $|A_{123}| = 1$ .

$$\begin{aligned} |A_1 \cup A_2 \cup A_3| &= |A_1| + |A_2| + |A_3| - |A_{12}| - |A_{13}| - |A_{23}| + |A_{123}| \\ &= 2 + 2 + 2 - 1 - 1 - 1 + 1 = 4. \end{aligned}$$


# Inclusion-Exclusion

$$|A_1 \cup A_2 \cup A_3| = |A_1| + |A_2| + |A_3| - |A_1 \cap A_2| - |A_1 \cap A_3| - |A_2 \cap A_3| + |A_1 \cap A_2 \cap A_3|.$$

*Proof.* Consider  $x \in A_2 \cap A_3$ . How many times is  $x$  counted?

$$\begin{array}{ccccccc} |A_1| & + & |A_2| & + & |A_3| & - & |A_1 \cap A_2| & - & |A_1 \cap A_3| & - & |A_2 \cap A_3| & + & |A_1 \cap A_2 \cap A_3| \\ 0 & & +1 & & +1 & & 0 & & 0 & & -1 & & 0 \end{array}$$

Contribution of  $x$  to sum is  $+1$ . Repeat for each region. ■



**Example (Derangements).** Give 3 coats to 3 girls so that noone gets their coat. How many ways?

$A_i = \{\text{girl } i \text{ gets her coat}\}$ .  $|A_i| = 2!$ .

$A_{ij} = \{\text{girls } i \text{ and } j \text{ get their coats}\}$ .  $|A_{ij}| = 1!$ .

$A_{123} = \{\text{girls 1, 2 and 3 get their coats}\}$ .  $|A_{123}| = 1$ .

$$\begin{aligned} |A_1 \cup A_2 \cup A_3| &= |A_1| + |A_2| + |A_3| - |A_{12}| - |A_{13}| - |A_{23}| + |A_{123}| \\ &= 2 + 2 + 2 - 1 - 1 - 1 + 1 = 4. \end{aligned}$$

The answer we seek is  $3! - 4 = 2$ .

(why?)

**Exercise.** How many numbers in  $1, \dots, 100$  are divisible by 2, 3 or 5?

# Pigeonhole Principle

---


If you have more guests than spare rooms, then some guests will have to share.

# Pigeonhole Principle

---

If you have more guests than spare rooms, then some guests will have to share.


- *More* pigeons than pigeonholes.
- A pigeonhole has two or more pigeons.



# Pigeonhole Principle

If you have more guests than spare rooms, then some guests will have to share.

- *More* pigeons than pigeonholes.
- A pigeonhole has two or more pigeons.



*Proof.* (By contraposition). Suppose no pigeonhole has 2 or more pigeons.

Let  $x_i$  be the number of pigeons in hole  $i$ ,  $x_i \leq 1$ .


$$\text{number of pigeons} = \sum_i x_i \leq \sum_i 1 = \text{number of pigeonholes.}$$

■

# Pigeonhole Principle

If you have more guests than spare rooms, then some guests will have to share.

- *More* pigeons than pigeonholes.
- A pigeonhole has two or more pigeons.



*Proof.* (By contraposition). Suppose no pigeonhole has 2 or more pigeons.

Let  $x_i$  be the number of pigeons in hole  $i$ ,  $x_i \leq 1$ .

$$\text{number of pigeons} = \sum_i x_i \leq \sum_i 1 = \text{number of pigeonholes.}$$


**Example.** If you have 8 people, at least two are born on the same day of the week.

We have 8 pigeons (the people) and 7 pigeonholes (the days of the week).

# Pigeonhole Principle

If you have more guests than spare rooms, then some guests will have to share.

- *More* pigeons than pigeonholes.
- A pigeonhole has two or more pigeons.



*Proof.* (By contraposition). Suppose no pigeonhole has 2 or more pigeons.

Let  $x_i$  be the number of pigeons in hole  $i$ ,  $x_i \leq 1$ .

$$\text{number of pigeons} = \sum_i x_i \leq \sum_i 1 = \text{number of pigeonholes.}$$

**Example.** If you have 8 people, at least two are born on the same day of the week.

We have 8 pigeons (the people) and 7 pigeonholes (the days of the week).

How many people do you need to ensure two are born on a Monday?

# Every Graph Has At Least One Pair of Social Twins

---

Two nodes are *social twins* if they have the same degree.





# Every Graph Has At Least One Pair of Social Twins

---

Two nodes are *social twins* if they have the same degree.



Assume the graph is connected.



# Every Graph Has At Least One Pair of Social Twins

Two nodes are *social twins* if they have the same degree.

Assume the graph is connected.



Degrees  $1, 2, \dots, (n - 1)$ , the pigeonholes. (Why no degree 0?)


Vertices  $v_1, v_2, \dots, v_n$ , the pigeons.

$n$  pigeons and  $(n - 1)$  pigeonholes, so at least two vertices are in the same degree-bin.

# Every Graph Has At Least One Pair of Social Twins

Two nodes are *social twins* if they have the same degree.

Assume the graph is connected.



Degrees  $1, 2, \dots, (n - 1)$ , the pigeonholes. (Why no degree 0?)


Vertices  $v_1, v_2, \dots, v_n$ , the pigeons.

$n$  pigeons and  $(n - 1)$  pigeonholes, so at least two vertices are in the same degree-bin.

If the graph is not connected, no one has degree  $n - 1$ .


Non-constructive proof: Who are those social twins? What are their degrees?

# Non-Constructive Proof and the Eye-Spy Dilemma






# Non-Constructive Proof and the Eye-Spy Dilemma



Prove to the 4 year old that the target exists in the picture

# Non-Constructive Proof and the Eye-Spy Dilemma


---



Prove to the 4 year old that the target exists in the picture



# Non-Constructive Proof and the Eye-Spy Dilemma



Prove to the 4 year old that the target exists in the picture

# Subset Sums

---

Given 100 twenty-seven digit numbers, find two subsets with the same subset-sum.

Prove that the professor is not sending the student on a wild-goose chase.



# Subset Sums

---

Given 100 twenty-seven digit numbers, find two subsets with the same subset-sum.

Prove that the professor is not sending the student on a wild-goose chase.

Any 27-digit number is at most  $10^{28}$ . So, a subset-sum is at most  $100 \times 10^{28} = 10^{30}$ .


# Subset Sums

---

Given 100 twenty-seven digit numbers, find two subsets with the same subset-sum.

Prove that the professor is not sending the student on a wild-goose chase.

Any 27-digit number is at most  $10^{28}$ . So, a subset-sum is at most  $100 \times 10^{28} = 10^{30}$ .



Pigeonholes: bins corresponding to each possible subset-sum,  $1, 2, \dots, 10^{30}$ .

Pigeons: the non-empty subsets of a 100-element set:  $2^{100} - 1 \approx 1.26 \times 10^{30}$  of them.


# Subset Sums

---

Given 100 twenty-seven digit numbers, find two subsets with the same subset-sum.

Prove that the professor is not sending the student on a wild-goose chase.

Any 27-digit number is at most  $10^{28}$ . So, a subset-sum is at most  $100 \times 10^{28} = 10^{30}$ .



Pigeonholes: bins corresponding to each possible subset-sum,  $1, 2, \dots, 10^{30}$ .

Pigeons: the non-empty subsets of a 100-element set:  $2^{100} - 1 \approx 1.26 \times 10^{30}$  of them.

At least two subsets must be in the same subset-sum-bin.

**Practice.** Exercise 14.6.

