

Foundations of Computer Science

Lecture 9

Sums And Asymptotics

Computing Sums

Asymptotics: big- $\Theta(\cdot)$, big- $O(\cdot)$, big- $\Omega(\cdot)$

The Integration Method

$$\sum_{k=1}^{\infty} \frac{(-1)^{k+1} k^2}{k^3 + 1} = ??$$



- ① Structural induction: proofs about recursively defined sets.
 - ▶ Matched parentheses.
 - ▶ $\mathbb{N}$
 - ▶ Palindromes.
 - ▶ Arithmetic expressions.
 - ▶ Rooted Binary Trees (RBT).

Today: Sums And Asymptotics

- 1 Maximum Substring Sum
- 2 Computing Sums
- 3 Asymptotics: Big-Theta, Big-Oh and Big-Omega
- 4 Integration Method

Maximum Substring Sum

1 -1 -1 2 3 4 -1 -1 2 3 -4 1 2 -1 -2 1

Maximum Substring Sum

1 -1 -1 2 3 4 -1 -1 2 3 -4 1 2 -1 -2 1

max. substring sum= 12

Maximum Substring Sum

1 -1 -1 2 3 4 -1 -1 2 3 -4 1 2 -1 -2 1

max. substring sum= 12

More generally, compute the maximum substring sum for

$$a_1 \quad a_2 \quad a_3 \quad a_4 \quad \cdots \quad a_{n-1} \quad a_n$$

Maximum Substring Sum

1 -1 -1 2 3 4 -1 -1 2 3 -4 1 2 -1 -2 1

max. substring sum= 12

More generally, compute the maximum substring sum for

$$a_1 \quad a_2 \quad a_3 \quad a_4 \quad \cdots \quad a_{n-1} \quad a_n$$

Different algorithms have different running times (n measures the “size” of the input),

$T_1(n) = 2 + \sum_{i=1}^n \left[2 + \sum_{j=i}^n \left(5 + \sum_{k=i}^j 2 \right) \right].$

(3 for loops)

$T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right).$

(2 for loops)

$T_3(n) = \begin{cases} 3 & n = 1; \\ 2T_3(\frac{1}{2}n) + 6n + 9 & n > 1 \text{ and even}; \\ T(\frac{1}{2}(n + 1)) + T(\frac{1}{2}(n - 1)) + 6n + 9 & n > 1 \text{ and odd.} \end{cases}$

(recursive)

$T_4(n) = 5 + \sum_{i=1}^n 10.$

(1 for loops)

(What does $\sum_{i=1}^n$ mean: Pop Quiz 9.1)

Maximum Substring Sum

1 -1 -1 2 3 4 -1 -1 2 3 -4 1 2 -1 -2 1

max. substring sum= 12

More generally, compute the maximum substring sum for

a_1 a_2 a_3 a_4 $\cdots$ a_{n-1} a_n

Different algorithms have different running times (n measures the “size” of the input),

$T_1(n) = 2 + \sum_{i=1}^n \left[2 + \sum_{j=i}^n \left(5 + \sum_{k=i}^j 2 \right) \right].$

(3 for loops)

$T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right).$

(2 for loops)

$T_3(n) = \begin{cases} 3 & n = 1; \\ 2T_3(\frac{1}{2}n) + 6n + 9 & n > 1 \text{ and even}; \\ T(\frac{1}{2}(n + 1)) + T(\frac{1}{2}(n - 1)) + 6n + 9 & n > 1 \text{ and odd.} \end{cases}$

(recursive)

$T_4(n) = 5 + \sum_{i=1}^n 10.$

(1 for loops)

(What does $\sum_{i=1}^n$ mean: Pop Quiz 9.1)

Which algorithm is best?

Evaluate the Runtimes

n	1	2	3	4	5	6	7	8	9	10
$T_1(n)$	11	29	58	100	157	231	324	438	575	737
$T_2(n)$	11	26	47	74	107	146	191	242	299	362

T_2 is better than T_1 ;

Evaluate the Runtimes

n	1	2	3	4	5	6	7	8	9	10
$T_1(n)$	11	29	58	100	157	231	324	438	575	737
$T_2(n)$	11	26	47	74	107	146	191	242	299	362
$T_3(n)$	3	27	57	87	123	159	195	231	273	315

T_2 is better than T_1 ;

T_2 versus T_3 ???

Evaluate the Runtimes

n	1	2	3	4	5	6	7	8	9	10
$T_1(n)$	11	29	58	100	157	231	324	438	575	737
$T_2(n)$	11	26	47	74	107	146	191	242	299	362
$T_3(n)$	3	27	57	87	123	159	195	231	273	315
$T_4(n)$	15	25	35	45	55	65	75	85	95	105

T_2 is better than T_1 ;

T_2 versus T_3 ???

What about T_4 ?

Evaluate the Runtimes

n	1	2	3	4	5	6	7	8	9	10
$T_1(n)$	11	29	58	100	157	231	324	438	575	737
$T_2(n)$	11	26	47	74	107	146	191	242	299	362
$T_3(n)$	3	27	57	87	123	159	195	231	273	315
$T_4(n)$	15	25	35	45	55	65	75	85	95	105

T_2 is better than T_1 ; T_2 versus T_3 ???

What about T_4 ?

We need:

- 1 Simple formulas for $T_1(n), \dots, T_4(n)$: we need to compute sums and solve recurrences.

Evaluate the Runtimes

n	1	2	3	4	5	6	7	8	9	10
$T_1(n)$	11	29	58	100	157	231	324	438	575	737
$T_2(n)$	11	26	47	74	107	146	191	242	299	362
$T_3(n)$	3	27	57	87	123	159	195	231	273	315
$T_4(n)$	15	25	35	45	55	65	75	85	95	105

T_2 is better than T_1 ; T_2 versus T_3 ???

What about T_4 ?

We need:

- 1 Simple formulas for $T_1(n), \dots, T_4(n)$: we need to compute sums and solve recurrences.
- 2 A way to compare runtime-*functions* that captures the essence of the algorithm.

Computing Sums: Tool 1: Constant Rule

$$S_1 = \sum_{i=1}^{10} 3$$

Computing Sums: Tool 1: Constant Rule

$$S_1 = \sum_{i=1}^{10} 3 = 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3$$

Computing Sums: Tool 1: Constant Rule

$$S_1 = \sum_{i=1}^{10} 3 = 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 \qquad 3 \times 10$$

Computing Sums: Tool 1: Constant Rule

$$S_1 = \sum_{i=1}^{10} 3 = 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 \qquad 3 \times 10$$

$$S_2 = \sum_{i=1}^{10} j$$

Computing Sums: Tool 1: Constant Rule

$$S_1 = \sum_{i=1}^{10} 3 = 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 \qquad 3 \times 10$$

$$S_2 = \sum_{i=1}^{10} j = j + j + j + j + j + j + j + j + j + j$$

Computing Sums: Tool 1: Constant Rule

$$S_1 = \sum_{i=1}^{10} 3 = 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 \qquad 3 \times 10$$

$$S_2 = \sum_{i=1}^{10} j = j + j + j + j + j + j + j + j + j + j \qquad j \times 10$$

Computing Sums: Tool 1: Constant Rule

$$S_1 = \sum_{i=1}^{10} 3 = 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 \qquad 3 \times 10$$

$$S_2 = \sum_{i=1}^{10} j = j + j + j + j + j + j + j + j + j + j \qquad j \times 10$$

$$S_3 = \sum_{i=1}^{10} i$$

Computing Sums: Tool 1: Constant Rule

$$S_1 = \sum_{i=1}^{10} 3 = 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 \qquad 3 \times 10$$

$$S_2 = \sum_{i=1}^{10} j = j + j + j + j + j + j + j + j + j + j \qquad j \times 10$$

$$S_3 = \sum_{i=1}^{10} i = 1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 + 9 + 10$$

Computing Sums: Tool 1: Constant Rule

$$S_1 = \sum_{i=1}^{10} 3 = 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 \qquad 3 \times 10$$

$$S_2 = \sum_{i=1}^{10} j = j + j + j + j + j + j + j + j + j + j \qquad j \times 10$$

$$S_3 = \sum_{i=1}^{10} i = 1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 + 9 + 10 \qquad \frac{1}{2} \times 10 \times (10 + 1)$$

Computing Sums: Tool 1: Constant Rule

$$S_1 = \sum_{i=1}^{10} 3 = 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 \qquad 3 \times 10$$

$$S_2 = \sum_{i=1}^{10} j = j + j + j + j + j + j + j + j + j + j \qquad j \times 10$$

$$S_3 = \sum_{i=1}^{10} i = 1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 + 9 + 10 \qquad \frac{1}{2} \times 10 \times (10 + 1)$$

The *index of summation* is i in these examples.

Computing Sums: Tool 1: Constant Rule

$$S_1 = \sum_{i=1}^{10} 3 = 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 \qquad 3 \times 10$$

$$S_2 = \sum_{i=1}^{10} j = j + j + j + j + j + j + j + j + j + j \qquad j \times 10$$

$$S_3 = \sum_{i=1}^{10} i = 1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 + 9 + 10 \qquad \frac{1}{2} \times 10 \times (10 + 1)$$

The *index of summation* is i in these examples.

Constants (independent of summation index) can be taken outside the sum.

$$S_1 = \sum_{i=1}^{10} 3 = 3 \sum_{i=1}^{10} 1 = 3 \times 10 \qquad S_2 = \sum_{i=1}^{10} j = j \sum_{i=1}^{10} 1 = j \times 10.$$

Computing Sums: Tool 1: Constant Rule

$$S_1 = \sum_{i=1}^{10} 3 = 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 + 3 \qquad 3 \times 10$$

$$S_2 = \sum_{i=1}^{10} j = j + j + j + j + j + j + j + j + j + j \qquad j \times 10$$

$$S_3 = \sum_{i=1}^{10} i = 1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 + 9 + 10 \qquad \frac{1}{2} \times 10 \times (10 + 1)$$

The *index of summation* is i in these examples.

Constants (independent of summation index) can be taken outside the sum.

$$S_1 = \sum_{i=1}^{10} 3 = 3 \sum_{i=1}^{10} 1 = 3 \times 10 \qquad S_2 = \sum_{i=1}^{10} j = j \sum_{i=1}^{10} 1 = j \times 10.$$

Pop Quiz 9.2 Compute $T_4(n) = 5 + \sum_{i=1}^n 10$.

Computing Sums: Tool 2: Addition Rule

$$S = \sum_{i=1}^5 (i + i^2)$$

Computing Sums: Tool 2: Addition Rule

$$\begin{aligned} S &= \sum_{i=1}^5 (i + i^2) \\ &= (1 + 1^2) + (2 + 2^2) + (3 + 3^2) + (4 + 4^2) + (5 + 5^2) \end{aligned}$$

Computing Sums: Tool 2: Addition Rule

$$\begin{aligned} S &= \sum_{i=1}^5 (i + i^2) \\ &= (1 + 1^2) + (2 + 2^2) + (3 + 3^2) + (4 + 4^2) + (5 + 5^2) \\ &= (1 + 2 + 3 + 4 + 5) + (1^2 + 2^2 + 3^2 + 4^2 + 5^2) \end{aligned} \quad \text{(rearrange terms)}$$

Computing Sums: Tool 2: Addition Rule

$$\begin{aligned} S &= \sum_{i=1}^5 (i + i^2) \\ &= (1 + 1^2) + (2 + 2^2) + (3 + 3^2) + (4 + 4^2) + (5 + 5^2) \\ &= (1 + 2 + 3 + 4 + 5) + (1^2 + 2^2 + 3^2 + 4^2 + 5^2) && \text{(rearrange terms)} \\ &= \sum_{i=1}^5 i + \sum_{i=1}^5 i^2. \end{aligned}$$

Computing Sums: Tool 2: Addition Rule

$$\begin{aligned} S &= \sum_{i=1}^5 (i + i^2) \\ &= (1 + 1^2) + (2 + 2^2) + (3 + 3^2) + (4 + 4^2) + (5 + 5^2) \\ &= (1 + 2 + 3 + 4 + 5) + (1^2 + 2^2 + 3^2 + 4^2 + 5^2) && \text{(rearrange terms)} \\ &= \sum_{i=1}^5 i + \sum_{i=1}^5 i^2. \end{aligned}$$

The sum of terms added together is the addition of the individual sums.

$$\sum_i (a(i) + b(i) + c(i) + \cdots) = \sum_i a(i) + \sum_i b(i) + \sum_i c(i) + \cdots$$

Computing Sums: Tool 3: Common Sums

$$\sum_{i=k}^n 1 = n + 1 - k$$

$$\sum_{i=1}^n i = \frac{1}{2}n(n+1)$$

$$\sum_{i=0}^n 2^i = 2^{n+1} - 1$$

$$\sum_{i=1}^n f(x) = nf(x)$$

$$\sum_{i=1}^n i^2 = \frac{1}{6}n(n+1)(2n+1)$$

$$\sum_{i=0}^n \frac{1}{2^i} = 2 - \frac{1}{2^n}$$

$$\sum_{i=0}^n r^i = \frac{1 - r^{n+1}}{1 - r} \quad (r \neq 1)$$

$$\sum_{i=1}^n i^3 = \frac{1}{4}n^2(n+1)^2$$

$$\sum_{i=1}^n \log i = \log n!$$

Example: $\sum_{i=1}^n (1 + 2i + 2^{i+2})$

$$\sum_{i=1}^n (1 + 2i + 2^{i+2}) =$$

Computing Sums: Tool 3: Common Sums

$$\sum_{i=k}^n 1 = n + 1 - k$$

$$\sum_{i=1}^n i = \frac{1}{2}n(n+1)$$

$$\sum_{i=0}^n 2^i = 2^{n+1} - 1$$

$$\sum_{i=1}^n f(x) = nf(x)$$

$$\sum_{i=1}^n i^2 = \frac{1}{6}n(n+1)(2n+1)$$

$$\sum_{i=0}^n \frac{1}{2^i} = 2 - \frac{1}{2^n}$$

$$\sum_{i=0}^n r^i = \frac{1 - r^{n+1}}{1 - r} \quad (r \neq 1)$$

$$\sum_{i=1}^n i^3 = \frac{1}{4}n^2(n+1)^2$$

$$\sum_{i=1}^n \log i = \log n!$$

Example: $\sum_{i=1}^n (1 + 2i + 2^{i+2})$

$$\sum_{i=1}^n (1 + 2i + 2^{i+2}) = \sum_{i=1}^n 1 + \sum_{i=1}^n 2i + \sum_{i=1}^n 2^{i+2} \quad (\text{addition rule})$$

Computing Sums: Tool 3: Common Sums

$$\sum_{i=k}^n 1 = n + 1 - k$$

$$\sum_{i=1}^n i = \frac{1}{2}n(n+1)$$

$$\sum_{i=0}^n 2^i = 2^{n+1} - 1$$

$$\sum_{i=1}^n f(x) = nf(x)$$

$$\sum_{i=1}^n i^2 = \frac{1}{6}n(n+1)(2n+1)$$

$$\sum_{i=0}^n \frac{1}{2^i} = 2 - \frac{1}{2^n}$$

$$\sum_{i=0}^n r^i = \frac{1 - r^{n+1}}{1 - r} \quad (r \neq 1)$$

$$\sum_{i=1}^n i^3 = \frac{1}{4}n^2(n+1)^2$$

$$\sum_{i=1}^n \log i = \log n!$$

Example: $\sum_{i=1}^n (1 + 2i + 2^{i+2})$

$$\sum_{i=1}^n (1 + 2i + 2^{i+2}) = \sum_{i=1}^n 1 + \sum_{i=1}^n 2i + \sum_{i=1}^n 2^{i+2} \quad (\text{addition rule})$$

$$= \sum_{i=1}^n 1 + 2 \sum_{i=1}^n i + 4 \sum_{i=1}^n 2^i \quad (\text{constant rule})$$

Computing Sums: Tool 3: Common Sums

$$\sum_{i=k}^n 1 = n + 1 - k$$

$$\sum_{i=1}^n i = \frac{1}{2}n(n+1)$$

$$\sum_{i=0}^n 2^i = 2^{n+1} - 1$$

$$\sum_{i=1}^n f(x) = nf(x)$$

$$\sum_{i=1}^n i^2 = \frac{1}{6}n(n+1)(2n+1)$$

$$\sum_{i=0}^n \frac{1}{2^i} = 2 - \frac{1}{2^n}$$

$$\sum_{i=0}^n r^i = \frac{1 - r^{n+1}}{1 - r} \quad (r \neq 1)$$

$$\sum_{i=1}^n i^3 = \frac{1}{4}n^2(n+1)^2$$

$$\sum_{i=1}^n \log i = \log n!$$

Example: $\sum_{i=1}^n (1 + 2i + 2^{i+2})$

$$\sum_{i=1}^n (1 + 2i + 2^{i+2}) = \sum_{i=1}^n 1 + \sum_{i=1}^n 2i + \sum_{i=1}^n 2^{i+2} \quad (\text{addition rule})$$

$$= \sum_{i=1}^n 1 + 2 \sum_{i=1}^n i + 4 \sum_{i=1}^n 2^i \quad (\text{constant rule})$$

$$= n + 2 \times \frac{1}{2}n(n+1) + 4 \cdot (2^{n+1} - 1 - \mathbf{1}) \quad (\text{common sums})$$

Computing Sums: Tool 3: Common Sums

$$\sum_{i=k}^n 1 = n + 1 - k$$

$$\sum_{i=1}^n i = \frac{1}{2}n(n+1)$$

$$\sum_{i=0}^n 2^i = 2^{n+1} - 1$$

$$\sum_{i=1}^n f(x) = nf(x)$$

$$\sum_{i=1}^n i^2 = \frac{1}{6}n(n+1)(2n+1)$$

$$\sum_{i=0}^n \frac{1}{2^i} = 2 - \frac{1}{2^n}$$

$$\sum_{i=0}^n r^i = \frac{1 - r^{n+1}}{1 - r} \quad (r \neq 1)$$

$$\sum_{i=1}^n i^3 = \frac{1}{4}n^2(n+1)^2$$

$$\sum_{i=1}^n \log i = \log n!$$

Example: $\sum_{i=1}^n (1 + 2i + 2^{i+2})$

$$\sum_{i=1}^n (1 + 2i + 2^{i+2}) = \sum_{i=1}^n 1 + \sum_{i=1}^n 2i + \sum_{i=1}^n 2^{i+2} \quad (\text{addition rule})$$

$$= \sum_{i=1}^n 1 + 2 \sum_{i=1}^n i + 4 \sum_{i=1}^n 2^i \quad (\text{constant rule})$$

$$= n + 2 \times \frac{1}{2}n(n+1) + 4 \cdot (2^{n+1} - 1 - 1) \quad (\text{common sums})$$

$$= n + n(n+1) + 2^{n+3} - 8 \quad (\text{common sums})$$

Computing Sums: Tool 3: Nested Sum Rule

$$S_1 = \sum_{i=1}^3 \sum_{j=1}^3 1;$$

$$S_2 = \sum_{i=1}^3 \sum_{j=1}^i 1.$$

Computing Sums: Tool 3: Nested Sum Rule

$$S_1 = \sum_{i=1}^3 \sum_{j=1}^3 1;$$

$$S_2 = \sum_{i=1}^3 \sum_{j=1}^i 1.$$

To compute a nested sum, start with the innermost sum and proceed outward.

Computing Sums: Tool 3: Nested Sum Rule

$$S_1 = \sum_{i=1}^3 \sum_{j=1}^3 1;$$

$$S_2 = \sum_{i=1}^3 \sum_{j=1}^i 1.$$

To compute a nested sum, start with the innermost sum and proceed outward.

$$S_1 \quad \sum_{\substack{j=1 \\ (i=1)}}^3 1 \quad \sum_{\substack{j=1 \\ (i=2)}}^3 1 \quad \sum_{\substack{j=1 \\ (i=3)}}^3 1$$

Computing Sums: Tool 3: Nested Sum Rule

$$S_1 = \sum_{i=1}^3 \sum_{j=1}^3 1;$$

$$S_2 = \sum_{i=1}^3 \sum_{j=1}^i 1.$$

To compute a nested sum, start with the innermost sum and proceed outward.

$$S_1 = \sum_{\substack{j=1 \\ (i=1)}}^3 1 + \sum_{\substack{j=1 \\ (i=2)}}^3 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1$$

Computing Sums: Tool 3: Nested Sum Rule

$$S_1 = \sum_{i=1}^3 \sum_{j=1}^3 1; \qquad S_2 = \sum_{i=1}^3 \sum_{j=1}^i 1.$$

To compute a nested sum, start with the innermost sum and proceed outward.

$$S_1 = \sum_{\substack{j=1 \\ (i=1)}}^3 1 + \sum_{\substack{j=1 \\ (i=2)}}^3 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1 = 3 + 3 + 3 = 9.$$

Computing Sums: Tool 3: Nested Sum Rule

$$S_1 = \sum_{i=1}^3 \sum_{j=1}^3 1;$$

$$S_2 = \sum_{i=1}^3 \sum_{j=1}^i 1.$$

To compute a nested sum, start with the innermost sum and proceed outward.

$$S_1 = \sum_{\substack{j=1 \\ (i=1)}}^3 1 + \sum_{\substack{j=1 \\ (i=2)}}^3 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1 = 3 + 3 + 3 = 9.$$

$$S_2 = \sum_{\substack{j=1 \\ (i=1)}}^1 1 + \sum_{\substack{j=1 \\ (i=2)}}^2 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1$$

Computing Sums: Tool 3: Nested Sum Rule

$$S_1 = \sum_{i=1}^3 \sum_{j=1}^3 1;$$

$$S_2 = \sum_{i=1}^3 \sum_{j=1}^i 1.$$

To compute a nested sum, start with the innermost sum and proceed outward.

$$S_1 = \sum_{\substack{j=1 \\ (i=1)}}^3 1 + \sum_{\substack{j=1 \\ (i=2)}}^3 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1 = 3 + 3 + 3 = 9.$$

$$S_2 = \sum_{\substack{j=1 \\ (i=1)}}^1 1 + \sum_{\substack{j=1 \\ (i=2)}}^2 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1$$

Computing Sums: Tool 3: Nested Sum Rule

$$S_1 = \sum_{i=1}^3 \sum_{j=1}^3 1;$$

$$S_2 = \sum_{i=1}^3 \sum_{j=1}^i 1.$$

To compute a nested sum, start with the innermost sum and proceed outward.

$$S_1 = \sum_{\substack{j=1 \\ (i=1)}}^3 1 + \sum_{\substack{j=1 \\ (i=2)}}^3 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1 = 3 + 3 + 3 = 9.$$

$$S_2 = \sum_{\substack{j=1 \\ (i=1)}}^1 1 + \sum_{\substack{j=1 \\ (i=2)}}^2 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1 = 1 + 2 + 3 = 6.$$

Computing Sums: Tool 3: Nested Sum Rule

$$S_1 = \sum_{i=1}^3 \sum_{j=1}^3 1;$$

$$S_2 = \sum_{i=1}^3 \sum_{j=1}^i 1.$$

To compute a nested sum, start with the innermost sum and proceed outward.

$$S_1 = \sum_{\substack{j=1 \\ (i=1)}}^3 1 + \sum_{\substack{j=1 \\ (i=2)}}^3 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1 = 3 + 3 + 3 = 9.$$

$$S_2 = \sum_{\substack{j=1 \\ (i=1)}}^1 1 + \sum_{\substack{j=1 \\ (i=2)}}^2 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1 = 1 + 2 + 3 = 6.$$

More generally:

$$S(n) = \sum_{i=1}^n \sum_{j=1}^i 1$$

Computing Sums: Tool 3: Nested Sum Rule

$$S_1 = \sum_{i=1}^3 \sum_{j=1}^3 1;$$

$$S_2 = \sum_{i=1}^3 \sum_{j=1}^i 1.$$

To compute a nested sum, start with the innermost sum and proceed outward.

$$S_1 = \sum_{\substack{j=1 \\ (i=1)}}^3 1 + \sum_{\substack{j=1 \\ (i=2)}}^3 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1 = 3 + 3 + 3 = 9.$$

$$S_2 = \sum_{\substack{j=1 \\ (i=1)}}^1 1 + \sum_{\substack{j=1 \\ (i=2)}}^2 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1 = 1 + 2 + 3 = 6.$$

More generally:

$$S(n) = \sum_{i=1}^n \sum_{j=1}^i 1 = \sum_{i=1}^n \underbrace{\sum_{j=1}^i 1}_{f(i)=i}$$

Computing Sums: Tool 3: Nested Sum Rule

$$S_1 = \sum_{i=1}^3 \sum_{j=1}^3 1;$$

$$S_2 = \sum_{i=1}^3 \sum_{j=1}^i 1.$$

To compute a nested sum, start with the innermost sum and proceed outward.

$$S_1 = \sum_{\substack{j=1 \\ (i=1)}}^3 1 + \sum_{\substack{j=1 \\ (i=2)}}^3 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1 = 3 + 3 + 3 = 9.$$

$$S_2 = \sum_{\substack{j=1 \\ (i=1)}}^1 1 + \sum_{\substack{j=1 \\ (i=2)}}^2 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1 = 1 + 2 + 3 = 6.$$

More generally:

$$S(n) = \sum_{i=1}^n \sum_{j=1}^i 1 = \sum_{i=1}^n \underbrace{\sum_{j=1}^i 1}_{f(i)=i} = \sum_{i=1}^n i$$

Computing Sums: Tool 3: Nested Sum Rule

$$S_1 = \sum_{i=1}^3 \sum_{j=1}^3 1;$$

$$S_2 = \sum_{i=1}^3 \sum_{j=1}^i 1.$$

To compute a nested sum, start with the innermost sum and proceed outward.

$$S_1 = \sum_{\substack{j=1 \\ (i=1)}}^3 1 + \sum_{\substack{j=1 \\ (i=2)}}^3 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1 = 3 + 3 + 3 = 9.$$

$$S_2 = \sum_{\substack{j=1 \\ (i=1)}}^1 1 + \sum_{\substack{j=1 \\ (i=2)}}^2 1 + \sum_{\substack{j=1 \\ (i=3)}}^3 1 = 1 + 2 + 3 = 6.$$

More generally:

$$S(n) = \sum_{i=1}^n \sum_{j=1}^i 1 = \sum_{i=1}^n \underbrace{\sum_{j=1}^i 1}_{f(i)=i} = \sum_{i=1}^n i = \frac{1}{2}n(n+1).$$

Computing a Formula for $T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right)$

$$T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right) =$$

Computing a Formula for $T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right)$

$$T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right) = 2 + \sum_{i=1}^n 3 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{sum rule})$$

Computing a Formula for $T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right)$

$$T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right) = 2 + \sum_{i=1}^n 3 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{sum rule})$$

$$= 2 + 3 \sum_{i=1}^n 1 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{constant rule})$$

Computing a Formula for $T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right)$

$$T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right) = 2 + \sum_{i=1}^n 3 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{sum rule})$$

$$= 2 + 3 \sum_{i=1}^n 1 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{constant rule})$$

$$= 2 + 3n + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{common sum})$$

Computing a Formula for $T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right)$

$$T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right) = 2 + \sum_{i=1}^n 3 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{sum rule})$$

$$= 2 + 3 \sum_{i=1}^n 1 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{constant rule})$$

$$= 2 + 3n + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{common sum})$$

$$= 2 + 3n + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{innermost sum})$$

Computing a Formula for $T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right)$

$$T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right) = 2 + \sum_{i=1}^n 3 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{sum rule})$$

$$= 2 + 3 \sum_{i=1}^n 1 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{constant rule})$$

$$= 2 + 3n + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{common sum})$$

$$= 2 + 3n + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{innermost sum})$$

$$= 2 + 3n + 6 \sum_{i=1}^n \sum_{j=i}^n 1 \quad (\text{constant rule})$$

Computing a Formula for $T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right)$

$$T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right) = 2 + \sum_{i=1}^n 3 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{sum rule})$$

$$= 2 + 3 \sum_{i=1}^n 1 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{constant rule})$$

$$= 2 + 3n + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{common sum})$$

$$= 2 + 3n + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{innermost sum})$$

$$= 2 + 3n + 6 \sum_{i=1}^n \sum_{j=i}^n 1 \quad (\text{constant rule})$$

$$= 2 + 3n + 6 \sum_{i=1}^n (n + 1 - i) \quad (\text{common sum})$$

Computing a Formula for $T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right)$

$$T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right) = 2 + \sum_{i=1}^n 3 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{sum rule})$$

$$= 2 + 3 \sum_{i=1}^n 1 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{constant rule})$$

$$= 2 + 3n + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{common sum})$$

$$= 2 + 3n + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{innermost sum})$$

$$= 2 + 3n + 6 \sum_{i=1}^n \sum_{j=i}^n 1 \quad (\text{constant rule})$$

$$= 2 + 3n + 6 \sum_{i=1}^n (n + 1 - i) \quad (\text{common sum})$$

$$= 2 + 3n + 6(n + (n - 1) + \cdots + 1)$$

Computing a Formula for $T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right)$

$$T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right) = 2 + \sum_{i=1}^n 3 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{sum rule})$$

$$= 2 + 3 \sum_{i=1}^n 1 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{constant rule})$$

$$= 2 + 3n + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{common sum})$$

$$= 2 + 3n + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{innermost sum})$$

$$= 2 + 3n + 6 \sum_{i=1}^n \sum_{j=i}^n 1 \quad (\text{constant rule})$$

$$= 2 + 3n + 6 \sum_{i=1}^n (n + 1 - i) \quad (\text{common sum})$$

$$= 2 + 3n + 6(n + (n - 1) + \cdots + 1)$$

$$= 2 + 3n + 6 \times \frac{1}{2}n(n + 1) \quad (\text{common sum})$$

Computing a Formula for $T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right)$

$$T_2(n) = 2 + \sum_{i=1}^n \left(3 + \sum_{j=i}^n 6 \right) = 2 + \sum_{i=1}^n 3 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{sum rule})$$

$$= 2 + 3 \sum_{i=1}^n 1 + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{constant rule})$$

$$= 2 + 3n + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{common sum})$$

$$= 2 + 3n + \sum_{i=1}^n \sum_{j=i}^n 6 \quad (\text{innermost sum})$$

$$= 2 + 3n + 6 \sum_{i=1}^n \sum_{j=i}^n 1 \quad (\text{constant rule})$$

$$= 2 + 3n + 6 \sum_{i=1}^n (n + 1 - i) \quad (\text{common sum})$$

$$= 2 + 3n + 6(n + (n - 1) + \cdots + 1)$$

$$= 2 + 3n + 6 \times \frac{1}{2}n(n + 1) \quad (\text{common sum})$$

$$= 2 + 6n + 3n^2 \quad (\text{algebra})$$

Practice: Compute a Formula for the Sum $\sum_{i=1}^n \sum_{j=1}^i ij$

$$\sum_{i=1}^n \sum_{j=1}^i ij =$$

Practice: Compute a Formula for the Sum $\sum_{i=1}^n \sum_{j=1}^i ij$

$$\sum_{i=1}^n \sum_{j=1}^i ij = \sum_{i=1}^n \sum_{j=1}^i ij \quad (\text{innermost sum})$$

Practice: Compute a Formula for the Sum $\sum_{i=1}^n \sum_{j=1}^i ij$

$$\begin{aligned} \sum_{i=1}^n \sum_{j=1}^i ij &= \sum_{i=1}^n \sum_{j=1}^i ij && \text{(innermost sum)} \\ &= \sum_{i=1}^n i \sum_{j=1}^i j && \text{(constant rule)} \end{aligned}$$

Practice: Compute a Formula for the Sum $\sum_{i=1}^n \sum_{j=1}^i ij$

$$\begin{aligned}\sum_{i=1}^n \sum_{j=1}^i ij &= \sum_{i=1}^n \sum_{j=1}^i ij && \text{(innermost sum)} \\ &= \sum_{i=1}^n i \sum_{j=1}^i j && \text{(constant rule)} \\ &= \sum_{i=1}^n i \times \frac{1}{2}i(i+1) && \text{(common sum)}\end{aligned}$$

Practice: Compute a Formula for the Sum $\sum_{i=1}^n \sum_{j=1}^i ij$

$$\sum_{i=1}^n \sum_{j=1}^i ij = \sum_{i=1}^n \sum_{j=1}^i ij \quad (\text{innermost sum})$$

$$= \sum_{i=1}^n i \sum_{j=1}^i j \quad (\text{constant rule})$$

$$= \sum_{i=1}^n i \times \frac{1}{2}i(i+1) \quad (\text{common sum})$$

$$= \frac{1}{2} \sum_{i=1}^n (i^3 + i^2) \quad (\text{algebra, constant rule})$$

Practice: Compute a Formula for the Sum $\sum_{i=1}^n \sum_{j=1}^i ij$

$$\sum_{i=1}^n \sum_{j=1}^i ij = \sum_{i=1}^n \sum_{j=1}^i ij \quad (\text{innermost sum})$$

$$= \sum_{i=1}^n i \sum_{j=1}^i j \quad (\text{constant rule})$$

$$= \sum_{i=1}^n i \times \frac{1}{2}i(i+1) \quad (\text{common sum})$$

$$= \frac{1}{2} \sum_{i=1}^n (i^3 + i^2) \quad (\text{algebra, constant rule})$$

$$= \frac{1}{2} \sum_{i=1}^n i^3 + \frac{1}{2} \sum_{i=1}^n i^2 \quad (\text{sum rule})$$

Practice: Compute a Formula for the Sum $\sum_{i=1}^n \sum_{j=1}^i ij$

$$\begin{aligned}\sum_{i=1}^n \sum_{j=1}^i ij &= \sum_{i=1}^n \sum_{j=1}^i ij && \text{(innermost sum)} \\ &= \sum_{i=1}^n i \sum_{j=1}^i j && \text{(constant rule)} \\ &= \sum_{i=1}^n i \times \frac{1}{2}i(i+1) && \text{(common sum)} \\ &= \frac{1}{2} \sum_{i=1}^n (i^3 + i^2) && \text{(algebra, constant rule)} \\ &= \frac{1}{2} \sum_{i=1}^n i^3 + \frac{1}{2} \sum_{i=1}^n i^2 && \text{(sum rule)} \\ &= \frac{1}{8}n^2(n+1)^2 + \frac{1}{12}n(n+1)(2n+1) && \text{(common sums)}\end{aligned}$$

Practice: Compute a Formula for the Sum $\sum_{i=1}^n \sum_{j=1}^i ij$

$$\begin{aligned}\sum_{i=1}^n \sum_{j=1}^i ij &= \sum_{i=1}^n \sum_{j=1}^i ij && \text{(innermost sum)} \\ &= \sum_{i=1}^n i \sum_{j=1}^i j && \text{(constant rule)} \\ &= \sum_{i=1}^n i \times \frac{1}{2}i(i+1) && \text{(common sum)} \\ &= \frac{1}{2} \sum_{i=1}^n (i^3 + i^2) && \text{(algebra, constant rule)} \\ &= \frac{1}{2} \sum_{i=1}^n i^3 + \frac{1}{2} \sum_{i=1}^n i^2 && \text{(sum rule)} \\ &= \frac{1}{8}n^2(n+1)^2 + \frac{1}{12}n(n+1)(2n+1) && \text{(common sums)} \\ &= \frac{1}{12}n + \frac{3}{8}n^2 + \frac{5}{12}n^3 + \frac{1}{8}n^4 && \text{(algebra)}\end{aligned}$$

Summary of Maximum Substring Sum Algorithms

Runtimes


$$\mathbf{T}_1(\mathbf{n}) = 2 + \frac{31}{6}n + \frac{7}{2}n^2 + \frac{1}{3}n^3$$

$$\mathbf{T}_2(\mathbf{n}) = 2 + 6n + 3n^2$$

$$3n(\log_2 n + 1) - 9 \leq \mathbf{T}_3(\mathbf{n}) \leq 12n(\log_2 n + 3) - 9$$

$$\mathbf{T}_4(\mathbf{n}) = 5 + 10n$$

(“simple” formulas for $T_1(n), \dots, T_4(n)$)



Summary of Maximum Substring Sum Algorithms

Runtimes


$$\mathbf{T_1(n)} = 2 + \frac{31}{6}n + \frac{7}{2}n^2 + \frac{1}{3}n^3$$

$$\mathbf{T_2(n)} = 2 + 6n + 3n^2$$

$$3n(\log_2 n + 1) - 9 \leq \mathbf{T_3(n)} \leq 12n(\log_2 n + 3) - 9$$

$$\mathbf{T_4(n)} = 5 + 10n$$

(“simple” formulas for $T_1(n), \dots, T_4(n)$)



So, which algorithm is best?

Summary of Maximum Substring Sum Algorithms

Runtimes


$$\mathbf{T_1(n)} = 2 + \frac{31}{6}n + \frac{7}{2}n^2 + \frac{1}{3}n^3$$

$$\mathbf{T_2(n)} = 2 + 6n + 3n^2$$

$$3n(\log_2 n + 1) - 9 \leq \mathbf{T_3(n)} \leq 12n(\log_2 n + 3) - 9$$

$$\mathbf{T_4(n)} = 5 + 10n$$

(“simple” formulas for $T_1(n), \dots, T_4(n)$)



So, which algorithm is best?

Computers solve problems with big inputs. We care about large n .

Summary of Maximum Substring Sum Algorithms

Runtimes


$$\mathbf{T_1(n)} = 2 + \frac{31}{6}n + \frac{7}{2}n^2 + \frac{1}{3}n^3$$

$$\mathbf{T_2(n)} = 2 + 6n + 3n^2$$

$$3n(\log_2 n + 1) - 9 \leq \mathbf{T_3(n)} \leq 12n(\log_2 n + 3) - 9$$

$$\mathbf{T_4(n)} = 5 + 10n$$

(“simple” formulas for $T_1(n), \dots, T_4(n)$)



So, which algorithm is best?

Computers solve problems with big inputs. We care about large n .

- Compare runtimes *asymptotically* in the input size n . That is $n \rightarrow \infty$.

Summary of Maximum Substring Sum Algorithms

Runtimes


$$\mathbf{T_1(n)} = 2 + \frac{31}{6}n + \frac{7}{2}n^2 + \frac{1}{3}n^3$$

$$\mathbf{T_2(n)} = 2 + 6n + 3n^2$$

$$3n(\log_2 n + 1) - 9 \leq \mathbf{T_3(n)} \leq 12n(\log_2 n + 3) - 9$$

$$\mathbf{T_4(n)} = 5 + 10n$$

(“simple” formulas for $T_1(n), \dots, T_4(n)$)



So, which algorithm is best?

Computers solve problems with big inputs. We care about large n .

- Compare runtimes *asymptotically* in the input size n . That is $n \rightarrow \infty$.
- Ignore additive and multiplicative constants (minutia). We care about *growth rate*.

Summary of Maximum Substring Sum Algorithms

Runtimes


$$\mathbf{T_1(n)} = 2 + \frac{31}{6}n + \frac{7}{2}n^2 + \frac{1}{3}n^3$$

$$\mathbf{T_2(n)} = 2 + 6n + 3n^2$$

$$3n(\log_2 n + 1) - 9 \leq \mathbf{T_3(n)} \leq 12n(\log_2 n + 3) - 9$$

$$\mathbf{T_4(n)} = 5 + 10n$$

(“simple” formulas for $T_1(n), \dots, T_4(n)$)



So, which algorithm is best?

Computers solve problems with big inputs. We care about large n .

- Compare runtimes *asymptotically* in the input size n . That is $n \rightarrow \infty$.
- Ignore additive and multiplicative constants (minutia). We care about *growth rate*.

Algorithm 4 is *linear* in n , $\frac{T_4(n)}{n} \rightarrow \text{constant}$.

Asymptotically Linear Functions: $\Theta(n)$, big-Theta-of- n

$T \in \Theta(\boldsymbol{n})$, if there are positive constants c, C for which

$$c \cdot \boldsymbol{n} \leq T(n) \leq C \cdot \boldsymbol{n}.$$

Asymptotically Linear Functions: $\Theta(n)$, big-Theta-of- n

$T \in \Theta(\mathbf{n})$, if there are positive constants c, C for which
 $c \cdot \mathbf{n} \leq T(n) \leq C \cdot \mathbf{n}$.

$$\frac{T(n)}{n} \xrightarrow{n \rightarrow \infty} \begin{cases} \infty & T \in \omega(n), & "T > n"; \\ \text{constant} > 0 & T \in \Theta(n), & "T = n"; \\ 0 & T \in o(n), & "T < n". \end{cases}$$

Asymptotically Linear Functions: $\Theta(n)$, big-Theta-of- n

$T \in \Theta(\mathbf{n})$, if there are positive constants c, C for which
 $c \cdot \mathbf{n} \leq T(n) \leq C \cdot \mathbf{n}$.

$$\frac{T(n)}{n} \xrightarrow{n \rightarrow \infty} \begin{cases} \infty & T \in \omega(n), & "T > n"; \\ \text{constant} > 0 & T \in \Theta(n), & "T = n"; \\ 0 & T \in o(n), & "T < n". \end{cases}$$

Linear means in $\Theta(n)$:

$$2n + 7, \quad 2n + 15\sqrt{n}, \quad 10^9 n + 3, \quad 3n + \log n, \quad 2^{\log_2 n + 4}.$$

Asymptotically Linear Functions: $\Theta(n)$, big-Theta-of- n

$T \in \Theta(\mathbf{n})$, if there are positive constants c, C for which
 $c \cdot \mathbf{n} \leq T(n) \leq C \cdot \mathbf{n}$.

$$\frac{T(n)}{n} \xrightarrow{n \rightarrow \infty} \begin{cases} \infty & T \in \omega(n), & "T > n"; \\ \text{constant} > 0 & T \in \Theta(n), & "T = n"; \\ 0 & T \in o(n), & "T < n". \end{cases}$$

Linear means in $\Theta(n)$:

$$2n + 7, \quad 2n + 15\sqrt{n}, \quad 10^9 n + 3, \quad 3n + \log n, \quad 2^{\log_2 n + 4}.$$

Not linear, not in $\Theta(n)$:

$$10^{-9}n^2, \quad 10^9\sqrt{n} + 15, \quad n^{1.0001}, \quad n^{0.9999}, \quad n \log n, \quad \frac{n}{\log n}, \quad 2^n.$$

Asymptotically Linear Functions: $\Theta(n)$, big-Theta-of- n

$$T \in \Theta(\mathbf{n}), \text{ if there are positive constants } c, C \text{ for which}$$
$$c \cdot \mathbf{n} \leq T(n) \leq C \cdot \mathbf{n}.$$

$$\frac{T(n)}{n} \xrightarrow{n \rightarrow \infty} \begin{cases} \infty & T \in \omega(n), & "T > n"; \\ \text{constant} > 0 & T \in \Theta(n), & "T = n"; \\ 0 & T \in o(n), & "T < n". \end{cases}$$

Linear means in $\Theta(n)$:

$$2n + 7, \quad 2n + 15\sqrt{n}, \quad 10^9n + 3, \quad 3n + \log n, \quad 2^{\log_2 n + 4}.$$

Not linear, not in $\Theta(n)$:


$$10^{-9}n^2, \quad 10^9\sqrt{n} + 15, \quad n^{1.0001}, \quad n^{0.9999}, \quad n \log n, \quad \frac{n}{\log n}, \quad 2^n.$$

Other runtimes from practice:

log	linear	loglinear	quadratic	cubic	superpolynomial	exponential	factorial	BAD
$\Theta(\log n)$	$\Theta(n)$	$\Theta(n \log n)$	$\Theta(n^2)$	$\Theta(n^3)$	$\Theta(n^{\log n})$	$\Theta(2^n)$	$\Theta(n!)$	$\Theta(n^n)$

General Asymptotics: $\Theta(f)$, big-Theta-of- f

$$\frac{T(n)}{f(n)} \xrightarrow{n \rightarrow \infty} \begin{cases} \infty & T \in \omega(f), \text{ “}T < f\text{”}; \\ \text{constant} > 0 & T \in \Theta(f), \text{ “}T = f\text{”}; \\ 0 & T \in o(f), \text{ “}T < f\text{”}. \end{cases}$$



$$T \in o(f)$$

$$\text{“}T < f\text{”}$$

$$T \in \Theta(f)$$

$$\text{“}T = f\text{”}$$

$$T \in \omega(f)$$


$$\text{“}T > f\text{”}$$

$$cf(n) \leq T(n) \leq Cf(n)$$

Examples and Practice. (See also Exercise 9.6)

General Asymptotics: $\Theta(f)$, big-Theta-of- f

$$\frac{T(n)}{f(n)} \xrightarrow{n \rightarrow \infty} \begin{cases} \infty & T \in \omega(f), \text{ “}T < f\text{”}; \\ \text{constant} > 0 & T \in \Theta(f), \text{ “}T = f\text{”}; \\ 0 & T \in o(f), \text{ “}T < f\text{”}. \end{cases}$$




$T \in o(f)$		$T \in \Theta(f)$		$T \in \omega(f)$
“ $T < f$ ”	“ $T \leq f$ ”	“ $T = f$ ”	“ $T \geq f$ ”	“ $T > f$ ”
$cf(n) \leq T(n) \leq Cf(n)$				

Examples and Practice. (See also Exercise 9.6)

General Asymptotics: $\Theta(f)$, big-Theta-of- f

$$\frac{T(n)}{f(n)} \xrightarrow{n \rightarrow \infty} \begin{cases} \infty & T \in \omega(f), \text{ “}T < f\text{”}; \\ \text{constant} > 0 & T \in \Theta(f), \text{ “}T = f\text{”}; \\ 0 & T \in o(f), \text{ “}T < f\text{”}. \end{cases}$$




$T \in o(f)$	$T \in O(f)$	$T \in \Theta(f)$	$T \in \Omega(f)$	$T \in \omega(f)$
“ $T < f$ ”	“ $T \leq f$ ”	“ $T = f$ ”	“ $T \geq f$ ”	“ $T > f$ ”
$cf(n) \leq T(n) \leq Cf(n)$				

Examples and Practice. (See also Exercise 9.6)

General Asymptotics: $\Theta(f)$, big-Theta-of- f

$$\frac{T(n)}{f(n)} \xrightarrow{n \rightarrow \infty} \begin{cases} \infty & T \in \omega(f), \text{ “}T < f\text{”}; \\ \text{constant} > 0 & T \in \Theta(f), \text{ “}T = f\text{”}; \\ 0 & T \in o(f), \text{ “}T < f\text{”}. \end{cases}$$




$T \in o(f)$	$T \in O(f)$	$T \in \Theta(f)$	$T \in \Omega(f)$	$T \in \omega(f)$
“ $T < f$ ”	“ $T \leq f$ ”	“ $T = f$ ”	“ $T \geq f$ ”	“ $T > f$ ”
	$T(n) \leq C f(n)$	$c f(n) \leq T(n) \leq C f(n)$	$c f(n) \leq T(n)$	

Examples and Practice. (See also Exercise 9.6)

General Asymptotics: $\Theta(f)$, big-Theta-of- f

$$\frac{T(n)}{f(n)} \xrightarrow{n \rightarrow \infty} \begin{cases} \infty & T \in \omega(f), \text{ “}T < f\text{”}; \\ \text{constant} > 0 & T \in \Theta(f), \text{ “}T = f\text{”}; \\ 0 & T \in o(f), \text{ “}T < f\text{”}. \end{cases}$$



$T \in o(f)$	$T \in O(f)$	$T \in \Theta(f)$	$T \in \Omega(f)$	$T \in \omega(f)$
“ $T < f$ ”	“ $T \leq f$ ”	“ $T = f$ ”	“ $T \geq f$ ”	“ $T > f$ ”
	$T(n) \leq C f(n)$	$c f(n) \leq T(n) \leq C f(n)$	$c f(n) \leq T(n)$	

Examples and Practice. (See also Exercise 9.6)


- For polynomials, growth rate is the highest order.

$$2n^2 \qquad n^2 + n\sqrt{n} \qquad n^2 + \log^{256} n \qquad n^2 + n^{1.99} \log^{256} n$$

$$\Theta(n^2) \qquad \Theta(n^2) \qquad \Theta(n^2) \qquad \Theta(n^2)$$

General Asymptotics: $\Theta(f)$, big-Theta-of- f

$$\frac{T(n)}{f(n)} \xrightarrow{n \rightarrow \infty} \begin{cases} \infty & T \in \omega(f), \text{ “}T < f\text{”}; \\ \text{constant} > 0 & T \in \Theta(f), \text{ “}T = f\text{”}; \\ 0 & T \in o(f), \text{ “}T < f\text{”}. \end{cases}$$




$T \in o(f)$	$T \in O(f)$	$T \in \Theta(f)$	$T \in \Omega(f)$	$T \in \omega(f)$
“ $T < f$ ”	“ $T \leq f$ ”	“ $T = f$ ”	“ $T \geq f$ ”	“ $T > f$ ”
	$T(n) \leq C f(n)$	$c f(n) \leq T(n) \leq C f(n)$	$c f(n) \leq T(n)$	

Examples and Practice. (See also Exercise 9.6)

- For polynomials, growth rate is the highest order.
- For nested sums, growth rate is number of nestings plus order of summand.

$2n^2$	$n^2 + n\sqrt{n}$	$n^2 + \log^{256} n$	$n^2 + n^{1.99} \log^{256} n$	$\sum_{i=1}^n i$	$\sum_{i=1}^n \sum_{j=1}^i 1$	$\sum_{i=1}^n \sum_{j=1}^i ij$
$\Theta(n^2)$	$\Theta(n^2)$	$\Theta(n^2)$	$\Theta(n^2)$	$\Theta(n^2)$	$\Theta(n^2)$	$\Theta(n^4)$


The Integration Method



Theorem. (Integration Bound)

For a monotonically increasing function f ,

The Integration Method




Theorem. (Integration Bound)

For a monotonically increasing function f ,


$$\int_{m-1}^n dx f(x) \leq \sum_{i=m}^n f(i)$$

The Integration Method

$$\int_0^n dx f(x)$$



$$\int_1^{n+1} dx f(x)$$




Theorem. (Integration Bound)

For a monotonically increasing function f ,


$$\int_{m-1}^n dx f(x) \leq \sum_{i=m}^n f(i)$$

The Integration Method

$$\int_0^n dx f(x)$$



$$\int_1^{n+1} dx f(x)$$




Theorem. (Integration Bound)

For a monotonically increasing function f ,


$$\int_{m-1}^n dx f(x) \leq \sum_{i=m}^n f(i) \leq \int_m^{n+1} dx f(x).$$

The Integration Method

$$\int_0^n dx f(x)$$



$$\int_1^{n+1} dx f(x)$$



Theorem. (Integration Bound)

For a monotonically increasing function f ,

$$\int_{m-1}^n dx f(x) \leq \sum_{i=m}^n f(i) \leq \int_m^{n+1} dx f(x).$$

(If f is monotonically decreasing, the inequalities are reversed.)

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k$$

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx x^k$$

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx \, x^k = \frac{n^{k+1}}{k+1}$$

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx \, x^k = \frac{n^{k+1}}{k+1} \in \Theta(n^{k+1}).$$

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx \, x^k = \frac{n^{k+1}}{k+1} \in \Theta(n^{k+1}).$$

- **Harmonic Numbers.** Set $f(x) = 1/x$ (monotonically decreasing):

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx x^k = \frac{n^{k+1}}{k+1} \in \Theta(n^{k+1}).$$

- **Harmonic Numbers.** Set $f(x) = 1/x$ (monotonically decreasing):

$$\underbrace{\int_1^{n+1} dx \frac{1}{x}}_{\ln(n+1)} \leq H_n = \sum_{i=1}^n \frac{1}{i}$$

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx x^k = \frac{n^{k+1}}{k+1} \in \Theta(n^{k+1}).$$

- **Harmonic Numbers.** Set $f(x) = 1/x$ (monotonically decreasing):

$$\underbrace{\int_1^{n+1} dx \frac{1}{x}}_{\ln(n+1)} \leq H_n = \sum_{i=1}^n \frac{1}{i} \leq \underbrace{1 + \int_1^n dx \frac{1}{x}}_{1+\ln n}.$$

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx x^k = \frac{n^{k+1}}{k+1} \in \Theta(n^{k+1}).$$

- **Harmonic Numbers.** Set $f(x) = 1/x$ (monotonically decreasing):

$$\underbrace{\int_1^{n+1} dx \frac{1}{x}}_{\ln(n+1)} \leq H_n = \sum_{i=1}^n \frac{1}{i} \leq \underbrace{1 + \int_1^n dx \frac{1}{x}}_{1+\ln n}.$$

- **Stirling's Approximation for $\ln n!$.** Set $f(x) = \ln x$:

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx x^k = \frac{n^{k+1}}{k+1} \in \Theta(n^{k+1}).$$

- **Harmonic Numbers.** Set $f(x) = 1/x$ (monotonically decreasing):

$$\underbrace{\int_1^{n+1} dx \frac{1}{x}}_{\ln(n+1)} \leq H_n = \sum_{i=1}^n \frac{1}{i} \leq \underbrace{1 + \int_1^n dx \frac{1}{x}}_{1+\ln n}.$$

- **Stirling's Approximation for $\ln n!$.** Set $f(x) = \ln x$:

$$\ln n! = \sum_{i=1}^n \ln i$$

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx x^k = \frac{n^{k+1}}{k+1} \in \Theta(n^{k+1}).$$

- **Harmonic Numbers.** Set $f(x) = 1/x$ (monotonically decreasing):

$$\underbrace{\int_1^{n+1} dx \frac{1}{x}}_{\ln(n+1)} \leq H_n = \sum_{i=1}^n \frac{1}{i} \leq \underbrace{1 + \int_1^n dx \frac{1}{x}}_{1+\ln n}.$$

- **Stirling's Approximation for $\ln n!$.** Set $f(x) = \ln x$:

$$\ln n! = \sum_{i=1}^n \ln i \leq \int_1^{n+1} dx \ln x$$

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx x^k = \frac{n^{k+1}}{k+1} \in \Theta(n^{k+1}).$$

- **Harmonic Numbers.** Set $f(x) = 1/x$ (monotonically decreasing):

$$\underbrace{\int_1^{n+1} dx \frac{1}{x}}_{\ln(n+1)} \leq H_n = \sum_{i=1}^n \frac{1}{i} \leq \underbrace{1 + \int_1^n dx \frac{1}{x}}_{1+\ln n}.$$

- **Stirling's Approximation for $\ln n!$.** Set $f(x) = \ln x$:

$$\ln n! = \sum_{i=1}^n \ln i \leq \int_1^{n+1} dx \ln x = (n+1) \ln(n+1) - n$$

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx x^k = \frac{n^{k+1}}{k+1} \in \Theta(n^{k+1}).$$

- **Harmonic Numbers.** Set $f(x) = 1/x$ (monotonically decreasing):

$$\underbrace{\int_1^{n+1} dx \frac{1}{x}}_{\ln(n+1)} \leq H_n = \sum_{i=1}^n \frac{1}{i} \leq \underbrace{1 + \int_1^n dx \frac{1}{x}}_{1+\ln n}.$$

- **Stirling's Approximation for $\ln n!$.** Set $f(x) = \ln x$:

$$\ln n! = \sum_{i=1}^n \ln i \leq \int_1^{n+1} dx \ln x = (n+1) \ln(n+1) - n \in \Theta(n \ln n).$$

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx x^k = \frac{n^{k+1}}{k+1} \in \Theta(n^{k+1}).$$

- **Harmonic Numbers.** Set $f(x) = 1/x$ (monotonically decreasing):

$$\underbrace{\int_1^{n+1} dx \frac{1}{x}}_{\ln(n+1)} \leq H_n = \sum_{i=1}^n \frac{1}{i} \leq \underbrace{1 + \int_1^n dx \frac{1}{x}}_{1+\ln n}.$$

- **Stirling's Approximation for $\ln n!$.** Set $f(x) = \ln x$:

$$\ln n! = \sum_{i=1}^n \ln i \leq \int_1^{n+1} dx \ln x = (n+1) \ln(n+1) - n \in \Theta(n \ln n).$$

- **Analyzing a Recurrence** $T_1 = 1; \quad T_n = T_{n-1} + n\sqrt{n} - \ln n.$

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx x^k = \frac{n^{k+1}}{k+1} \in \Theta(n^{k+1}).$$

- **Harmonic Numbers.** Set $f(x) = 1/x$ (monotonically decreasing):

$$\underbrace{\int_1^{n+1} dx \frac{1}{x}}_{\ln(n+1)} \leq H_n = \sum_{i=1}^n \frac{1}{i} \leq \underbrace{1 + \int_1^n dx \frac{1}{x}}_{1+\ln n}.$$

- **Stirling's Approximation for $\ln n!$.** Set $f(x) = \ln x$:

$$\ln n! = \sum_{i=1}^n \ln i \leq \int_1^{n+1} dx \ln x = (n+1) \ln(n+1) - n \in \Theta(n \ln n).$$

- **Analyzing a Recurrence** $T_1 = 1; \quad T_n = T_{n-1} + n\sqrt{n} - \ln n.$

First unfold the recurrence

$$T_n = \cancel{T_{n-1}} + n\sqrt{n} - \ln n$$

$$\cancel{T_{n-1}} = \cancel{T_{n-2}} + (n-1)\sqrt{n-1} - \ln(n-1)$$

$\vdots$

$$\cancel{T_3} = \cancel{T_2} + 3\sqrt{3} - \ln 3$$

$$\cancel{T_2} = \cancel{T_1}^1 + 2\sqrt{2} - \ln 2$$

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx x^k = \frac{n^{k+1}}{k+1} \in \Theta(n^{k+1}).$$

- **Harmonic Numbers.** Set $f(x) = 1/x$ (monotonically decreasing):

$$\underbrace{\int_1^{n+1} dx \frac{1}{x}}_{\ln(n+1)} \leq H_n = \sum_{i=1}^n \frac{1}{i} \leq \underbrace{1 + \int_1^n dx \frac{1}{x}}_{1+\ln n}.$$

- **Stirling's Approximation for $\ln n!$.** Set $f(x) = \ln x$:

$$\ln n! = \sum_{i=1}^n \ln i \leq \int_1^{n+1} dx \ln x = (n+1) \ln(n+1) - n \in \Theta(n \ln n).$$

- **Analyzing a Recurrence** $T_1 = 1; \quad T_n = T_{n-1} + n\sqrt{n} - \ln n.$

First unfold the recurrence

$$T_n = \cancel{T_{n-1}} + n\sqrt{n} - \ln n$$

$$\cancel{T_{n-1}} = \cancel{T_{n-2}} + (n-1)\sqrt{n-1} - \ln(n-1)$$

$\vdots$

$$\cancel{T_3} = \cancel{T_2} + 3\sqrt{3} - \ln 3$$

$$\cancel{T_2} = \cancel{T_1}^1 + 2\sqrt{2} - \ln 2$$

$$+ T_n = 1 + 2\sqrt{2} + \cdots + n\sqrt{n} - (\ln 2 + \ln 3 + \cdots + \ln n)$$

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx x^k = \frac{n^{k+1}}{k+1} \in \Theta(n^{k+1}).$$

- **Harmonic Numbers.** Set $f(x) = 1/x$ (monotonically decreasing):

$$\underbrace{\int_1^{n+1} dx \frac{1}{x}}_{\ln(n+1)} \leq H_n = \sum_{i=1}^n \frac{1}{i} \leq \underbrace{1 + \int_1^n dx \frac{1}{x}}_{1+\ln n}.$$

- **Stirling's Approximation for $\ln n!$.** Set $f(x) = \ln x$:

$$\ln n! = \sum_{i=1}^n \ln i \leq \int_1^{n+1} dx \ln x = (n+1) \ln(n+1) - n \in \Theta(n \ln n).$$

- **Analyzing a Recurrence** $T_1 = 1; \quad T_n = T_{n-1} + n\sqrt{n} - \ln n.$

First unfold the recurrence

$$T_n = \cancel{T_{n-1}} + n\sqrt{n} - \ln n$$

$$\cancel{T_{n-1}} = \cancel{T_{n-2}} + (n-1)\sqrt{n-1} - \ln(n-1)$$

$\vdots$

$$\cancel{T_3} = \cancel{T_2} + 3\sqrt{3} - \ln 3$$

$$\cancel{T_2} = \cancel{T_1}^1 + 2\sqrt{2} - \ln 2$$

$$+ T_n = 1 + 2\sqrt{2} + \cdots + n\sqrt{n} - (\ln 2 + \ln 3 + \cdots + \ln n)$$

$$= \sum_{i=1}^n i\sqrt{i} - \sum_{i=1}^n \ln i$$

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx x^k = \frac{n^{k+1}}{k+1} \in \Theta(n^{k+1}).$$

- **Harmonic Numbers.** Set $f(x) = 1/x$ (monotonically decreasing):

$$\underbrace{\int_1^{n+1} dx \frac{1}{x}}_{\ln(n+1)} \leq H_n = \sum_{i=1}^n \frac{1}{i} \leq \underbrace{1 + \int_1^n dx \frac{1}{x}}_{1+\ln n}.$$

- **Stirling's Approximation for $\ln n!$.** Set $f(x) = \ln x$:

$$\ln n! = \sum_{i=1}^n \ln i \leq \int_1^{n+1} dx \ln x = (n+1) \ln(n+1) - n \in \Theta(n \ln n).$$

- **Analyzing a Recurrence** $T_1 = 1; \quad T_n = T_{n-1} + n\sqrt{n} - \ln n.$

First unfold the recurrence

$$\begin{aligned} T_n &= \cancel{T_{n-1}} + n\sqrt{n} - \ln n \\ \cancel{T_{n-1}} &= \cancel{T_{n-2}} + (n-1)\sqrt{n-1} - \ln(n-1) \\ &\vdots \end{aligned}$$

$$\cancel{T_3} = \cancel{T_2} + 3\sqrt{3} - \ln 3$$

$$\cancel{T_2} = \cancel{T_1} + 2\sqrt{2} - \ln 2$$

$$+ T_n = 1 + 2\sqrt{2} + \cdots + n\sqrt{n} - (\ln 2 + \ln 3 + \cdots + \ln n)$$

$$= \sum_{i=1}^n i\sqrt{i} - \sum_{i=1}^n \ln i$$

$$T(n) = \underbrace{\sum_{i=1}^n i\sqrt{i}}_{\Theta(n^{5/2})} - \underbrace{\sum_{i=1}^n \ln i}_{\ln n! \in \Theta(n \ln n)}$$

Integration For Quickly Getting Asymptotic Behavior

- **Integer Powers.** Set $f(x) = x^k$:

$$\sum_{i=1}^n i^k \approx \int_0^n dx \, x^k = \frac{n^{k+1}}{k+1} \in \Theta(n^{k+1}).$$

- **Harmonic Numbers.** Set $f(x) = 1/x$ (monotonically decreasing):

$$\underbrace{\int_1^{n+1} dx \, \frac{1}{x}}_{\ln(n+1)} \leq H_n = \sum_{i=1}^n \frac{1}{i} \leq \underbrace{1 + \int_1^n dx \, \frac{1}{x}}_{1+\ln n}.$$

- **Stirling's Approximation for $\ln n!$.** Set $f(x) = \ln x$:

$$\ln n! = \sum_{i=1}^n \ln i \leq \int_1^{n+1} dx \, \ln x = (n+1) \ln(n+1) - n \in \Theta(n \ln n).$$

- **Analyzing a Recurrence** $T_1 = 1; \quad T_n = T_{n-1} + n\sqrt{n} - \ln n.$

First unfold the recurrence

$$\begin{aligned} T_n &= \cancel{T_{n-1}} + n\sqrt{n} - \ln n \\ \cancel{T_{n-1}} &= \cancel{T_{n-2}} + (n-1)\sqrt{n-1} - \ln(n-1) \\ &\vdots \\ \cancel{T_3} &= \cancel{T_2} + 3\sqrt{3} - \ln 3 \\ \cancel{T_2} &= \cancel{T_1} + 2\sqrt{2} - \ln 2 \\ \hline + T_n &= 1 + 2\sqrt{2} + \cdots + n\sqrt{n} - (\ln 2 + \ln 3 + \cdots + \ln n) \\ &= \sum_{i=1}^n i\sqrt{i} - \sum_{i=1}^n \ln i \end{aligned}$$

$$T(n) = \underbrace{\sum_{i=1}^n i\sqrt{i}}_{\Theta(n^{5/2})} - \underbrace{\sum_{i=1}^n \ln i}_{\ln n! \in \Theta(n \ln n)}$$

$$T(n) \in \Theta(n^{5/2}).$$

