An Analysis of the Maximum Drawdown Risk Measure

Malik Magdon-Ismail (RPI)

November 14, 2005.

Joint work with:

Amir Atiya (Cairo University)
Amrit Pratap(Caltech)
Yaser Abu-Mostafa(Caltech)

Example

Fund	$\mu(\%)$	$\sigma(\%)$	max. DD(%)	T (yrs)
Π_1	25	10	-5	1
Π_2	30	10	$-7\frac{1}{2}$	1.5
Π_3	25	12.5	$-8\frac{1}{3}$	2

$$Calmar\ Ratio = \frac{\text{Return over } [0,T]}{\text{max. DD over } [0,T]}$$

$$Sterling\ Ratio = \frac{\text{Return over } [0,T]}{\text{max. DD over } [0,T] - 10\%}$$

Which fund is best?

The Complication

- μ and σ are annualized.
- Max. drawdown is computed over a given time period.
- **Problem**: funds have statistics over different length time intervals.

How do we annualize MDD?

Why?

Common Practice

Compare funds over $T = 3$ yrs.

Artificial:

- Wasteful of useful data.
- 3 years data may not be available.
- Easily available data does not generally quote the MDD for 3 years.

$\sqrt{T}$-Rule for Sharpe Ratio

$$Sharpe\ Ratio = \frac{\mu(\text{annualized})}{\sigma(\text{annualized})}$$

$$\left.\begin{array}{l}\mu_\tau \\ \sigma_\tau\end{array}\right\}\text{over time periods of size } \tau$$

$$\begin{aligned}\mu(\text{annualized}) &= \mu_\tau \cdot \frac{1}{\tau} \\ \sigma(\text{annualized}) &= \sigma_\tau \cdot \sqrt{\frac{1}{\tau}}\end{aligned}$$

$$\boxed{Sharpe\ Ratio = \frac{\mu_\tau}{\sigma_\tau \sqrt{\tau}}}$$

Similar scaling laws for Sterling-type Ratios?

Part I:

Analysis of the Maximum Drawdown.

Part II:

Application to Scaling Laws.

Part I:

Analysis of Maximum Drawdown

The Drawdown (DD)

The DD (current loss) is the loss from the peak to the current value.

$X(t)$ is the cumulative return curve.



$$DD(T) = \sup_{s \in [0,T]} X(s) - X(T)$$

$DD(T)$ is well understood (eg. **[Karatzas and Shreve, 1997]**).

The Maximum Drawdown (MDD)

The MDD is the maximum loss incurred from a peak to a bottom.



$$MDD(T) = \sup_{t \in [0,T]} DD(t)$$

– DD is an extremum.
– MDD is an **extremum of an extremum**.

Sampling of Prior Work

– Previous work on the the MDD is mostly empirical or Monte-Carlo.

- [Acar and James; 1997]
- [Sornette; 2002]
- [Burghardt, Duncan and Liu; 2003]
- [Harding, Nakou and Nejjar; 2003]
- [Chekhlov, Uryasev and Zabarankin; 2003]

– The only analytical approach is for a Brownian motion with **zero drift**, [Douady, Shiryaev and Yor; 2000].

Setup

$X(t)$ is an (arithmetic) Brownian motion:

$$\boxed{\boldsymbol{dX(t) = \mu dt + \sigma dW(t) \qquad\qquad 0 \leq t \leq T}}$$

$\mu =$	average return per unit time	(drift)
$\sigma =$	std. dev. of the returns per unit time	(volatility)
$dW(t) =$	Wiener increment	(shocks)

Note: If the fund $S(t)$ follows a geometric Brownian motion, then the cumulative return sequence follows a Brownian motion.

We would like to study $MDD(\mu, \sigma, T)$.

$DD(t)$ is a Reflected Brownian Motion

Drawdown at time t is a stochastic process.



$DD(t)$ is reflecting at 0, i.e., DD(t) is a **reflected Brownian motion**.

$$dDD(t) = \begin{cases} -dX(t) & DD(t) > 0 \\ \max\{0, -dX(t)\} & DD(t) = 0. \end{cases}$$

[Magdon-Ismail, Atiya, Pratap and Abu-Mostafa; 2004]

Expected $MDD(\mu, \sigma, T)$

Theorem: $$E[MDD(\mu,\sigma,T)] = \begin{cases} \frac{2\sigma^2}{\mu} Q_p\left(\frac{\mu^2 T}{2\sigma^2}\right) & \mu > 0 \\ \sqrt{\frac{\pi}{2}}\sigma\sqrt{T} & \mu = 0 \\ \frac{2\sigma^2}{\mu} Q_n\left(\frac{\mu^2 T}{2\sigma^2}\right) & \mu < 0 \end{cases}$$

Q_p and Q_n are **"universal" functions**.

– Only need to be computed once!

– Asymptotics?

[Magdon-Ismail, Atiya, Pratap and Abu-Mostafa; 2004]

Dimensionless quantity: $x = \dfrac{\mu\sqrt{T}}{\sigma}$.

Behavior of $Q_{MDD}(x)$



http://www.cs.rpi.edu/~magdon/data/Qfunctions.html

Part II:

Scaling Laws

Recap

$$\text{Calmar}(T) = \frac{\text{Return over } [0,T]}{MDD \text{ over } [0,T]} \approx \frac{\mu T}{E[MDD]} = \text{Clmr}$$

Given two funds,

$\Pi_1 : \mu_1, \sigma_1, T_1, MDD_1, \text{Clmr}_1 = \frac{\mu_1 T_1}{MDD_1}$.

$\Pi_2 : \mu_2, \sigma_2, T_2, MDD_2, \text{Clmr}_2 = \frac{\mu_2 T_2}{MDD_2}$.

How to compare Clmr_1 and Clmr_2?

Normalized Calmar Ratio

Normalize the ratios to a reference time τ, for example $\tau = 1$ yr.

We know how to scale return over $[0, T_1]$,

$$\mu_1 T_1 \longrightarrow \mu_1 \tau = \mu_1 T_1 \cdot \frac{\tau}{T_1}$$

We can scale $MDD([0, T_1]) \to MDD([0, \tau])$ using proportion,

$$\frac{E[MDD([0, \tau])]}{E[MDD([0, T_1])]} = \frac{MDD([0, \tau])}{MDD([0, T_1])}$$

Example – Revisited

Fund	$\mu(\%)$	$\sigma(\%)$	max. DD(%)	T (yrs)	$\mathbf{Calmar}$	$\overline{\mathbf{Calmar}}$
Π_1	25	10	-5	1	**5**	**5**
Π_2	30	10	$-7\frac{1}{2}$	1.5	**6**	**4.41**
Π_3	25	12.5	$-8\frac{1}{3}$	2	**6**	**3.62**

(normalized to $\tau = 1$ yr.)

$$\boxed{\Pi_1 > \Pi_2 > \Pi_3}$$

The Relative Strength β

Consider the long horizon, $\tau \to \infty$:

$$\beta = \frac{\overline{Calmar}}{\overline{Calmar}_{\text{ref.}}},$$

(eg. reference instrument = S&P 500).

β defines a total order.

Example – Re-Revisited

Fund	$\mu(\%)$	$\sigma(\%)$	max. DD(%)	T (yrs)	β
Π_1	25	10	-5	1	**1.00**
Π_2	30	10	$-7\frac{1}{2}$	1.5	**0.97**
Π_3	25	12.5	$-8\frac{1}{3}$	2	**0.64**

(relative strengths w.r.t Π_1.)

$$\boxed{\Pi_1 > \Pi_2 > \Pi_3}$$

Real Data*

Fund	μ(%)	σ(%)	T(yrs)	MDD	$Calmar$	**E[MDD]**	$\overline{\textbf{Calmar}}$	$\boldsymbol{\beta}$
$S\&P500$	10.04	15.48	24.25	46.28	5.261	**44.56**	**0.6104**	**1**
$FTSE100$	7.01	16.66	19.83	48.52	2.865	**55.54**	**0.4395**	**0.5003**
$NASDAQ$	11.20	24.38	19.42	75.04	2.899	**77.87**	**0.4402**	**0.5407**
DCM	15.65	5.78	3.08	3.11	15.50	**4.770**	**6.541**	**27.76**
NLT	3.35	16.03	3.08	25.40	0.4062	**31.35**	**0.2202**	**0.1331**
OIC	17.19	4.52	1.16	0.42	47.48	**2.493**	**42.31**	**212.0**
TGF	8.48	9.83	4.58	8.11	4.789	**15.84**	**1.752**	**3.589**

DCM=Diamond Capital Management;
NLT=Non-Linear Technologies;
OIC=Olsen Investment Corporation;
TGF=Tradewinds Global Fund.

– Normalized Calmar ratio is to $\tau = 1$ yr.

– Relative strength index β is computed w.r.t. $S\&P500$.

*International Advisory Services Group http://iasg.pertrac2000.com/mainframe.asp

Conclusion

1. Studied MDD for a Brownian motion.

2. We now have scaling laws for MDD and Sterling-type ratios.

3. Can compare trading strategies over different time intervals.

Advertisement:

[Magdon-Ismail, Atiya, Pratap, Abu-Mostafa; 2004]

[Magdon-Ismail, Atiya; 2004]

http://www.cs.rpi.edu/~magdon

Thank You

References

[1] **I. Karatzas and S. E. Shreve.** *Brownian Motion and Stochastic Calculus*. Springer, 1997.

[2] **E. Acar and S. James**, *Maximum loss and maximum drawdown in financial markets*. Int. Conference on Forecasting Financial Markets, London, UK, 1997.

[3] **D. Sornette**, *Why Do Stock Markets Crash: Critical Events in Complex Financial Systems*. Princeton University Press, 2002.

[4] **Burghardt, G., Duncan, R. and L. Liu**, *Deciphering drawdown*. Risk magazine, Risk management for investors, September, S16-S20, 2003.

[5] **D. Harding, G. Nakou, and A. Nejjar**, *The pros and cons of drawdown as a statistical measure for risk in investments*. AIMA Journal, pp. 16-17, April 2003.

[6] **Chekhlov, A., Uryasev, S., and M. Zabarankin**, *Drawdown Measure in Portfolio Optimization*. Research Report 2003-15. ISE Dept., University of Florida, September 2003.

[7] **R. Douady, A. Shiryaev, and M. Yor**, *On the probability characteristics of downfalls in a standard Brownian motion*. SIAM, Theory Probability Appl., Vol 44, pp. 29-38, 2000.

[8] **M. Dominé** *First passage time distribution of a Wiener process with drift concerning two elastic barries*. Journal of Applied Probability, DD:164–175, 1996.

[9] **M. Magdon-Ismail, A. Atiya, A. Pratap and Y. Abu-Mostafa**, *On the Maximum Drawdown of a Brownian Motion*. Journal of Applied Probability, Volume 41, Number 1, 2004.

[10] **M. Magdon-Ismail, A. Atiya**, *Maximum Drawdown*. Risk Magazine, Volume 17, Number 10, pages 99-102, October, 2004.